

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of which leaves are the organs, and to consider it as a merely unavoidable evaporation whose amount depends upon physical factors that need to be exactly evaluated. What would be thought of an engineer who attempted to compare the performance of two boilers of different construction if he knew nothing of the heat units applied? Yet most of the so-called "comparative" studies of transpiration take no account of the fundamental energetics involved, assuming that when leaves are placed "under the same conditions" they have a like amount of energy for transpiration!—C. R. B.

Correlation and leaf size.—A. J. EWART has published a short note dealing with correlation and leaf size.<sup>28</sup> LINDEMUTH has shown that adult leaves of Begonia and Iresine increase in size when allowed to root in the soil. EWART conducted experiments on *Tilia europaea*, and found that adult leaves do not increase in size, even when most of the leaves are removed. However, a partial defoliation of young shoots causes a development of unusually large leaves, and the increase in size beyond the normal is due to an increase in the number of cells and not to an increase in their size, as supposed by LINDEMUTH. That increased size is due to an increased number of cells in such cases is not the common view.—H. C. COWLES.

Presynapsis and synapsis.—Presynaptic and synaptic stages in the first division of the embryo sac mother cell of Adoxa are interpreted by LAGERBERG as follows.<sup>29</sup> Before synapsis the chromatin granules collect into groups which are often in pairs, apparently less numerous than the mature chromosomes. The chromatic substance of the groups becomes distributed along parallel linin threads, so that at the beginning of synapsis there are two parallel threads which fuse as synapsis proceeds. The single thread which is thus formed is the thickest which appears at any time in this nucleus. Accordingly, the fusion of the male and female elements of the nucleus takes place during synapsis.—Charles J. Chamberlain.

Embryology of Capsella.—The late Mrs. Mabel Schaffner<sup>30</sup> left an incomplete paper on the embryology of Capsella, which her husband, Professor John H. Schaffner, has published. It is a detailed study of the development of the embryo of what is perhaps the most frequently used dicotyledon in teaching. Aside from the completeness of the series, the striking feature of the plates is that the whole series is drawn to the same scale, representing to the eye the actual increase of the embryo in size at each stage of its development.—J. M. C.

<sup>&</sup>lt;sup>28</sup> EWART, A. J., The influence of correlation upon the size of leaves. Annals of Botany 20:79-82. 1906.

<sup>&</sup>lt;sup>29</sup> LAGERBERG, TORSTEN, Ueber die präsynaptische und synaptische Entwickelung der Kerne in der Embryosackmutterzellen von *Adaxa moschatellina*. Botaniska Studier, tillägnade F. R. KJELLMAN. 1906:80–88.

<sup>30</sup> Schaffner, Mabel, The embryology of the shepherd's purse. Ohio Nat. 7:1-8. pls. 1-3. 1906.